

Cheap Nursing CEUs

Fragile X Syndrome

1. Fragile X syndrome is more common in females.

- A. True
 - B. False
-

2. Which of the following may be considered a physical symptom associated with Fragile X syndrome?

- A. Large head
 - B. Large teeth
 - C. Small ears
 - D. Wide face
-

3. Which of the following may be considered a behavioral, social, and emotional symptom associated with Fragile X syndrome?

- A. Hyperactivity
 - B. An inability to learn new skills
 - C. An inability to efficiently or effectively solve problems
 - D. Stuttering
-

4. Which of the following statements is most accurate?

- A. Often individuals show symptoms of Fragile X syndrome by 2 months to 4 months of age.
 - B. Often individuals show symptoms of Fragile X syndrome by 6 months to 10 months of age.
 - C. The physical symptoms associated with Fragile X syndrome typically develop when an individual is about 6 months of age.
 - D. The physical symptoms associated with Fragile X syndrome typically develop when an individual goes through puberty.
-

5. Which of the following may be considered an early sign/symptom of Fragile X syndrome?

- A. Begins to speak early
- B. Learns to walk early
- C. Displays unusual reactions to the way things sound, smell, taste, look, or feel

D. Enlarged testicles in males

6. In order to be diagnosed with an intellectual disability, an individual must have both a significantly low IQ and considerable problems in everyday functioning.

- A. True**
 - B. False**
-

7. A patient has questions regarding Celexa. Which of the following educational points should be expressed to the patient?

- A. Celexa is a SSRI.**
 - B. Celexa is an antipsychotic agent.**
 - C. Celexa is not associated with nausea or dry mouth.**
 - D. A typical dose of Celexa is 100 mg daily.**
-

8. A patient has questions regarding Concerta. Which of the following educational points should be expressed to the patient?

- A. Concerta is a SSRI.**
 - B. Concerta is an antipsychotic agent.**
 - C. A typical starting dose of Concerta for children and adolescents is 80 mg once daily.**
 - D. A typical starting dose of Concerta for children and adolescents is 18 mg once daily.**
-

9. What is a typical dose of Adderall XR for individuals ages 6 - 17?

- A. 10 mg once daily**
 - B. 50 mg once daily**
 - C. 100 mg once daily**
 - D. 100 mg twice daily**
-

10. A genetic counselor is an individual who specializes in genetics and provides patients with counseling, education, and assistance in making decisions regarding genetic-related health care.

- A. True**
 - B. False**
-